

Improved approximations of Poissonian errors for high confidence levels

Harald Ebeling

Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

2 February 2008

ABSTRACT

We present improved numerical approximations to the exact Poissonian confidence limits for small numbers n of observed events following the approach of Gehrels (1986). Analytic descriptions of all parameters used in the approximations are provided to allow their straightforward inclusion in computer algorithms for processing of large data sets. Our estimates of the upper (lower) Poisson confidence limits are accurate to better than 1% for $n \leq 100$ and values of S , the derived significance in units of Gaussian standard deviations, of up to 7 (5). In view of the slow convergence of the commonly used Gaussian approximations toward the correct Poissonian values, in particular for higher values of S , we argue that, for $n \leq 40$, Poissonian statistics should be used in most applications, unless errors of the order of, or exceeding, 10% are acceptable.

Key words: methods: statistical — methods: numerical

1 INTRODUCTION

The need to assess the statistical significance of an observed small number of events is common in astrophysics as well as in virtually all other natural sciences. The number of neutrinos detected in an underground detector, the number of supernovae observed at $z > 1$, the number of photons in a faint X-ray point source — all of these are numbers likely to be in the Poisson regime, and knowing the correct errors of these measurements is obviously crucial to any scientific conclusions drawn from these numbers.

In a much noted paper Gehrels (1986, from here on G86) provided analytic approximations to the correct Poisson confidence limits for small event numbers. A particularly useful new and improved approximation in G86 is for the Poisson lower confidence limit, λ_l (see Section 2 below for an overview of the nomenclature used). However, two issues limit the applicability of these approximations to large data sets and/or applications that require high confidence levels. First, Gehrels’ analytic approximation to the correct value of λ_l uses empirically determined parameters β and γ , both of which are non-trivial functions of S , the desired significance in units of Gaussian standard deviations. G86 tabulates $\beta(S)$ and $\gamma(S)$ for ten values of S ranging from one to 3.3σ but does not provide an analytic formula that would allow the reader to compute λ_l for arbitrary S values¹. Second, the validity of all approximations presented and discussed in G86 has only been verified in the same, relatively narrow range of confidence levels from 1 to 3.3σ . To our knowledge, no reasonably accurate approximations have been published for $S > 3.3\sigma$.

While these issues may be of limited importance for many, if not most, applications, they can become serious in cases where confidence limits need to be computed for large sets of event numbers, particularly if high confidence levels are required. Consider, for instance, X-ray astronomy, a traditionally photon starved line of observational research. With large-area, high-resolution X-ray CCD detectors now in use on board the Chandra and XMM-Newton X-ray Observatories, X-ray images of dimensions 1024×1024 or even 4096×4096 pixels have become common. The numbers of photons registered in the vast majority of these pixels will be in the Poisson regime, and assessing accurately the significance of any features embedded in the very low and spatially non-uniform background measured with these large arrays requires the accurate computation of a considerable number of Poisson confidence limits. A real-life example of a scientific project relevant in this context is the compilation of a statistical sample of unresolved (single-pixel) point sources detected at the greater than 5σ confidence level (i.e., $S = 5$) in a set of Chandra ACIS-I images.

It is for applications like the ones outlined above that we here present modified and improved versions of G86’s approximations that are more accurate over an extended range of confidence levels ($S \lesssim 7\sigma$). To permit these approximations to be incorporated straightforwardly into computer algorithms for the processing of large data sets, we also provide analytic descriptions of all parameters used, thus allowing the computation of accurate Poisson confidence limits for a wide range of event numbers and confidence levels.

¹ Especially relevant in view of the fact that $\gamma(S)$ actually becomes singular close to $S = 1$, as we show in Section 5.

2 NOMENCLATURE

In the following we adopt Gehrels' nomenclature and definitions. Specifically, upper limits λ_u and lower limits λ_l are, for Poisson statistics, defined by

$$\sum_{x=0}^n \frac{\lambda_u^n e^{-\lambda_u}}{x!} = 1 - \text{CL} \quad n \geq 0 \quad (1)$$

and

$$\sum_{x=0}^{n-1} \frac{\lambda_l^n e^{-\lambda_l}}{x!} = \text{CL} \quad n \geq 1 \quad (2)$$

where n is the number of events observed, and CL is the desired confidence level (Eq. 1 and 2 of G86). (For $n = 0$, $\lambda_l = 0$ for all values of CL). For Gaussian statistics (i.e. a probability distribution which is normal) CL is related to S , the equivalent Gaussian number of σ , by

$$\text{CL}(S) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^S e^{-t^2/2} dt. \quad (3)$$

For ease of presentation we shall use S to parametrize a large range of CL values.

For $n = 0$ to 100 and selected values of S ranging from 1 to 3.3 G86 tabulates λ_u and λ_l as obtained from Eqs. 1 and 2; Gehrels also presents analytic and numerical approximations accurate to better than 2% for $S < 3.3$. In this paper, we test the Gaussian approximation as well as the ones discussed in G86 over a larger range of confidence levels ($S \leq 7$). We then modify Gehrels approximations to improve their performance specifically for large values of S and, finally, present numerical (polynomial) descriptions of all parameters used, to allow the computation of approximate values of $\lambda_u(n, S)$ and $\lambda_l(n, S)$ for all values of n up to at least 100 and $S \leq 7$.

3 THE GAUSSIAN APPROXIMATION

For a normal probability distribution (i.e., in the limit $n \rightarrow \infty$) λ_u and λ_l are given by the well known expressions $n \pm S\sqrt{n}$ which are also commonly used to approximate the not straightforwardly computable Poisson limits for 'reasonably' large values of n . What 'reasonable' means in this context is subject to debate; in practice, values of n in excess of 20 (and sometimes even $n > 10$) are often deemed sufficiently large to justify the use of the Gaussian approximation.

In Figure 1, we show the percentage error of the Gaussian approximation $n \pm S\sqrt{n}$ when applied to event numbers in the Poisson regime (loosely defined as $n < 100$). For low to moderate confidence levels ($S \leq 2$) the Gaussian approximation is accurate to better than 10% for $n = 20$ (but not for $n = 10$!), which may or may not be sufficient for a given application. However, the error of the Gaussian approximation, in particular for the lower limit, increases rapidly for higher confidence levels. Already at $S = 3$, $n > 45$ is required to limit the error in λ_l to 10%; for $S = 5$ even $n = 100$ is insufficient if 10% accuracy are sought. 10% accuracy may not nearly be good enough though in applications where errors of many independent measurements are propagated. In the case of X-ray spectral fitting, for instance, theoretical models are fitted simultaneously to events registered in hundreds of independent energy channels. A *systematic* error of the order of 10% in the errors

on the counts in *all* spectral bins introduced by the use of Gaussian approximations to the true Poisson errors can lead to best-fit parameter values that may be erroneous by significantly more than the formal, statistical errors.

4 APPROXIMATION OF THE POISSONIAN UPPER LIMIT

G86 derives two analytic approximations to the true Poissonian upper limit, namely

$$\lambda_u \approx (n+1) \left[1 - \frac{1}{9(n+1)} + \frac{S}{3\sqrt{n+1}} \right]^3 \quad (\text{G86 Eq. 9})$$

and

$$\lambda_u \approx n + S\sqrt{n+1} + \frac{S^2+2}{3} \quad (\text{G86 Eq. 10})$$

where the latter expression is simply the previous one expanded and limited to the dominant terms in $(n+1)$.

In Figure 2 we show the percentage error of these two approximations for a range of n and S values. Both expressions represent a considerable improvement over the Gaussian approximations (cf. Fig. 1). At low values of n the remaining error of several per cent may, however, still be too high for certain applications. Following the approach taken by Gehrels to improve his approximation to λ_l (see the following Section) we therefore apply a heuristic correction to G86 Eq. 9 in the form of an additional term $b(n+1)^c$:

$$\lambda_u \approx (n+1) \left[1 - \frac{1}{9(n+1)} + \frac{S}{3\sqrt{n+1}} + b(n+1)^c \right]^3. \quad (4)$$

We determine $b = b(S)$ from the requirement that the above approximation be an identity for $n = 0$, i.e.,

$$b(S) = \lambda_u(0, S)^{1/3} - 1 + 1/9 - S/3, \quad (5)$$

thus forcing better performance for low values of n . For each value of S from 0 to 7, $c = c(S)$ is then chosen such that the error of the above approximation is minimized for $0 \leq n \leq 100$. The runs of b and c as functions of S are shown in Figure 3. We find c to be negative for all values of S with one-sided singularities at the roots of Eq. 5, which lie at $S_{0,1} = 0.50688$ (corresponding to a confidence level of 69.388%) and $S_{0,2} = 2.27532$ (confidence level 98.856%).

To allow the evaluation of Eq. 4 for any number of observed events, n , and any confidence level, S , specifically in the proximity of the mentioned singularities, we fit piecewise polynomial functions to $b(S)$ and $c(1/S)$, or $c(\log_{10} S)$, with the degree of the polynomial being determined by the requirement that the absolute of the residuals be less than 1% over the full S range of the fit, except at the locations of the mentioned singularities. We find acceptable polynomial descriptions of $b(S)$ and $c(S)$ as follows:

$$b(S) = \sum_{i=0}^8 b_i S^i \quad (6)$$

and

$$c(S) = \begin{cases} \sum_{i=0}^4 c_{1,i} \left(\frac{1}{S-S_{0,1}} \right)^i & : S < S_{0,1} \\ \sum_{i=0}^4 c_{2,i} \log_{10}(S-S_{0,1})^i & : S_{0,1} < S < 1.2 \\ \sum_{i=0}^3 c_{3,i} \left(\frac{1}{S-S_{0,2}} \right)^i & : 1.2 < S < S_{0,2} \\ \sum_{i=0}^7 c_{4,i} \log_{10}(S-S_{0,2})^i & : S > S_{0,2} \end{cases} \quad (7)$$

with $-10 \leq c(S) \leq 0$ overriding the above definition where necessary, and coefficients $b_i, c_{1,i}, c_{2,i}, c_{3,i}$, and $c_{4,i}$ as listed in Table 1. The results of these fits are shown as the solid lines in Fig. 3.

Figure 4 shows the relative, absolute errors of Eq. 4 and demonstrates that our approximation is accurate to better than 0.5% for all values of n and S considered here.

5 APPROXIMATION OF THE POISSONIAN LOWER LIMIT

As evidenced by Fig. 1 the Gaussian approximation $\lambda_l(n, S) \approx n - S\sqrt{n}$ is a poor one for all but the lowest values of S . G86 explores the behaviour of several more sophisticated analytic approximations before resorting to modifying the most promising of them by adding a heuristic power law term βn^γ (we used the same approach in the preceding section to improve the approximation to λ_u):

$$\lambda_l \approx n \left(1 - \frac{1}{9n} + \frac{S}{3\sqrt{n}} + \beta n^\gamma \right)^3. \quad (\text{G86 Eq. 14})$$

To find $\beta(S)$ and $\gamma(S)$ we proceed similarly as before for our approximation to λ_u and define $\beta(S)$ as

$$\beta(S) = \lambda_l(1, S)^{1/3} - 1 + 1/9 + S/3, \quad (8)$$

and then determine $\gamma(S)$ such that the error of the above approximation is minimized for $0 \leq n \leq 100$. The result, β and γ as functions of S , is shown in Figure 5 which, in the overlap region, agrees with Fig. 1 of G86. γ is negative for all values of S with a one-sided singularity at the only root of Eq. 8 at $S_0 = 0.93876$, corresponding to a confidence level of 82.607%.

To facilitate the evaluation of Eq. 14 of G86 for a wide range of values of n and S , we attempt to find analytical expressions for $\beta(S)$ and $\gamma(S)$ (G86 quotes the values of either function only at 10 locations between $S = 1$ and $S = 3.291$). In analogy to the approach taken in the preceding section, we fit piecewise polynomial functions to $\beta(S)$, as well as to $\gamma(S)$, $\gamma(1/S)$, or $\gamma(\log_{10} S)$, with the degree of the polynomial being determined by the requirement that the absolute of the residuals be less than 2% over the full S range of the fit (less than 0.1% at the high- S end where high accuracy is critical). We find acceptable polynomial descriptions of $\beta(S)$ and $\gamma(S)$ as follows:

$$\beta(S) = \begin{cases} \sum_{i=0}^5 \beta_{1,i} S^i & : S \leq 3 \\ \sum_{i=0}^4 \beta_{2,i} S^i & : S > 3 \end{cases} \quad (9)$$

and

$$\gamma(S) = \begin{cases} \sum_{i=0}^5 \gamma_{1,i} \log_{10}(S_0 - S)^i & : S \leq S_0 \\ \sum_{i=0}^5 \gamma_{2,i} \left(\frac{1}{S-S_0}\right)^i & : S_0 < S \leq 2.7 \\ \sum_{i=0}^2 \gamma_{3,i} S^i & : S > 2.7 \end{cases} \quad (10)$$

with $-50 \leq \gamma(S) \leq 0$ overriding the above where necessary, and coefficients $\beta_{1,i}, \beta_{2,i}, \gamma_{1,i}, \gamma_{2,i}$, and $\gamma_{3,i}$ as listed in Table 2. The results of these fits are shown as the solid lines in Fig. 5.

Equation 14 of G86 indeed yields greatly reduced errors when compared to the exact values of λ_l . For $n < 100$ and $1 < S < 3.291$ G86 quotes an accuracy of better than 2% for the above approximation. Figure 6 confirms this, but also demonstrates that the errors become unacceptably large ($> 10\%$) for higher confidence levels and small to moderate values of n .

We now attempt to improve on Eq. 14 of G86 by adding a second, higher-order correction term. As illustrated by Fig. 6 such

an additional term would have to improve the performance of the approximation particularly in the high- S regime for which Eq. 14 of G86 was not optimized. This goal can be achieved by introducing a (totally ad-hoc) sinusoidal term which adds only one additional parameter δ :

$$\lambda_l \approx n \left[1 - \frac{1}{9n} + \frac{S}{3\sqrt{n}} + \beta n^\gamma + \delta \sin \left(\frac{5}{n+1/4} \frac{\pi}{2} \right) \right]^3. \quad (11)$$

Since the sinusoidal term, by design, vanishes for $n = 1$, Eq. 8 still holds, and continues to define $\beta(S)$. $\gamma(S)$ (slightly different from the one determined from Eq. 14 of G86) and $\delta(S)$ are again obtained by iteratively minimizing the absolute error of the approximation for $0 \leq n \leq 100$. With $\beta(S)$ unchanged and $\gamma(S)$ virtually indistinguishable from the data shown in Fig. 5 we can focus on $\delta(S)$ which shows a complex behaviour (Fig. 7). We do not attempt to model the run of $\delta(S)$ for small values of S where the function remains close to zero. Instead, we fit a high-order polynomial to the high- S end and set $\delta(S) = 0$ for $S < 1.2$:

$$\delta(S) = \begin{cases} 0 & : S < 1.2 \\ \sum_{i=0}^8 \delta_i S^i & : S \geq 1.2 \end{cases} \quad (12)$$

with coefficients δ_i as listed in Table 3. The results of these fits are shown as the solid line in Fig. 7.

Figure 8 demonstrates that Eq. 11 provides an approximation to λ_l that is accurate to better than 1% when the polynomial fits to $\beta(S)$, $\gamma(S)$, and $\delta(S)$ (Eqs. 9, 10, 12) are used, except for $n = 1$ where an error of just over 1% is observed.

6 SUMMARY

The Gaussian approximation $\lambda_u \approx n + S\sqrt{n}$ to the true Poissonian upper confidence limit is acceptable for low confidence levels ($S < 3\sigma$) and $n > 40$, but becomes increasingly inaccurate for higher values of S . The situation is worse for the Gaussian approximation $\lambda_l \approx n - S\sqrt{n}$ to the true Poissonian lower confidence limit, which is off by more than 10% at $S = 5$ even at $n = 100$. The approximations proposed by G86 greatly improve upon the Gaussian estimates but are still inaccurate at the 10% level for low values of n and high confidence levels.

Building on Gehrels' work we present improved algebraic approximations which reduce the error with respect to the true Poissonian confidence limits to under 1% for $S \leq 7$ (Poisson upper limit) and $S \leq 5$ (Poisson lower limit). Although we have tested these equations only for $n \leq 100$, their analytic behaviour suggests that they hold for all values of n (cf. Figures 4 and 8).

To allow the numerical computation of approximate Poissonian confidence limits for arbitrary combinations of n and S within the quoted ranges, we provide the coefficients of piecewise polynomial fits to all parameters used in the definition of either approximation.

All figures of this paper were produced using the Interactive Data Language (IDL); the IDL source code of the approximations `poisson_uplim` (Eq. 4) and `poisson_lolim` (Eq. 11) is available from the author.

HE gratefully acknowledges financial support from NASA LTSA grant NAG 5-8253 and NASA ADP grant NAG 5-9238.

4 Harald Ebeling

i	b_i	$c_{1,i}$	$c_{2,i}$	$c_{3,i}$	$c_{4,i}$
0	-3.8954e-03	-2.0799e+00	-1.4354e+00	-8.4098e-01	-1.0120e+00
1	+6.2328e-03	-7.1925e-01	-6.3188e-01	+6.8766e-01	-2.8853e-01
2	+5.2345e-03	-4.0064e-01	-1.6177e-01	+2.0358e-01	+4.2013e-01
3	-5.3096e-03	-7.3386e-02	-5.6966e-01	+3.9965e-02	-5.3310e-02
4	+1.3093e-03	-5.4791e-03	-2.2835e-01		-1.6319e-02
5	-2.0344e-04				+4.8667e-02
6	+2.0393e-05				-5.5299e-02
7	-1.1974e-06				-3.3361e-02
8	+3.1161e-08				

Table 1. Coefficients of polynomial fits to $b(S)$ and $c(S)$ of Eq. 4, as defined in Eqs. 6 and 7

i	$\beta_{1,i}$	$\beta_{2,i}$	$\gamma_{1,i}$	$\gamma_{2,i}$	$\gamma_{3,i}$
0	-3.8605809e-03	+3.4867327e-01	-1.7480435	-0.6347351	-2.7517416e+00
1	-6.6002964e-03	-4.0996949e-01	-1.8895824	-4.6707845	+3.1692400e-01
2	+6.5798149e-03	+1.6514495e-01	-3.0808786	+6.1602866	-8.7788310e-03
3	+2.8172041e-03	-1.5783156e-02	-5.5164953	-4.3543401	
4	+2.9892915e-03	+5.2768918e-04	-3.9940504	+1.4470675	
5	-5.4387574e-04		-1.0248451	-0.1870896	

Table 2. Coefficients of polynomial fits to $\beta(S)$ and $\gamma(S)$ of G86, Eq. 14, as defined in Eqs. 9 and 10

REFERENCES

Gehrels N. 1986, ApJ, 303, 336 (G86)

i	$\beta_{1,i}$	$\beta_{2,i}$	$\gamma_{1,i}$	$\gamma_{2,i}$	$\gamma_{3,i}$	δ_i
0	-3.8605809e-03	+3.4867327e-01	-1.7174713	-1.0131243	-2.8115538e+00	-2.2906640e-02
1	-6.6002964e-03	-4.0996949e-01	-1.7015942	-2.9319339	3.5117552e-01	+6.8209168e-02
2	+6.5798149e-03	+1.6514495e-01	-1.9059468	+3.2459998	-1.3215426e-02	-9.1678422e-02
3	+2.8172041e-03	-1.5783156e-02	-3.1324250	-2.1348935		+7.1533924e-02
4	+2.9892915e-03	+5.2768918e-04	-2.0145052	+0.6676902		-3.5010270e-02
5	-5.4387574e-04		-0.4257810	-0.0834041		+1.0928872e-02
6						-2.1069241e-03
7						+2.2638722e-04
8						-1.0302360e-05

Table 3. Coefficients of polynomial fits to $\beta(S)$, $\gamma(S)$, and $\delta(S)$ of Eq. 11, as defined in Eqs. 9, 10, and 12




Figure 1. Percentage error of the Gaussian approximations $\lambda_u \approx n + S\sqrt{n}$ (top) and $\lambda_l \approx n - S\sqrt{n}$ (bottom) as a function of n . In each panel the S values of the shown curves vary from lower left to upper right in steps of 0.5 as indicated; the thick lines correspond to $S = 1, 2$, and 3 . Note how, for $n = 10$ (marked by the dotted line), the errors still reach and exceed 10% for all but the lowest values of S .




Figure 2. Percentage error of the approximations of G86 Eq. 9 (top) and G86 Eq. 10 (bottom) as a function of n . In each panel the S values of the shown curves vary from 1 to 7 as annotated. For essentially all values of n and S explored here the error of either approximation remains below the 10% level for $n > 2$, and below 1% for $n \gtrsim 35$.




Figure 3. Run of parameters b (top) and c (bottom) of Eq. 4 as a function of S , the equivalent number of Gaussian σ . c exhibits singularities at $S = 0.507$ and $S = 2.275$ where $b = 0$. The bullets mark the locations at which $b(S)$ and $c(S)$ were computed, the solid lines mark polynomial fits to the data (see text for details).




Figure 4. Percentage error of the approximations of Eq. 4 with $b(S)$ and $c(S)$ as computed (top), and using the polynomial fits of Eqs. 6 and 7 (bottom), as a function of n . In each panel the S values of the shown curves vary from 0.5 to 7 as annotated. For all values of n and S explored here the error remains below the 0.5% level.




Figure 5. Run of parameters β (top) and γ (bottom) of G86 Eq. 14 as a function of S , the equivalent number of Gaussian σ . γ exhibits a singularity at $S = 0.939$ where $\beta = 0$. The bullets mark the locations at which $\beta(S)$ and $\gamma(S)$ were computed, the solid lines mark polynomial fits to the data (see text for details).




Figure 6. Percentage error of the approximation to λ_l given by G86 Eq. 14 with $\beta(S)$ and $\gamma(S)$ as computed (top), and using the polynomial fits of Eqs. 9 and 10 (bottom), as a function of n . In each panel the S values of the shown curves vary from 1 to 5 as annotated. While the approximation is good for low to moderate confidence levels ($S < 3.5$) it fails for $S > 4$ where errors approaching and exceeding 10% are observed for small values of n .




Figure 7. Run of δ of Eq. 11 as a function of S , the equivalent number of Gaussian σ . The bullets mark the locations at which $\delta(S)$ was computed, the solid line marks a polynomial fit to the data (see text for details).




Figure 8. Percentage error of the approximations of Eq. 11 with $\beta(S)$, $\gamma(S)$, and $\delta(S)$ as computed (top), and using the polynomial fits of Eqs. 9, 10, and 12 (bottom) as a function of n . In each panel the S values of the shown curves vary from 0.5 to 5 as annotated. For all values of n (except $n = 1$) and S explored here the error remains below the 1% level.